

Definitions and examples of Finsler spacetimes

MIGUEL ANGEL JAVALOYES VICTORIA

*Department of Mathematics, University of Murcia
majava@um.es*

May 8, 2017

ABSTRACT

We will analyze the different definitions of Finsler spacetimes provided by Beem [3], Asanov [2], Pfeifer-Wohlfarth [7], Kostelecky [6], Javaloyes-Sánchez [4], Amir-Azami [1], describing their advantages and disadvantages in every case, and what type of results can be obtained. Finally we will describe how to generate a big quantity of examples in order to test the theory [5].

References

- [1] A. B. AAZAMI AND M. A. JAVALOYES, *Penrose's singularity theorem in a Finsler spacetime*, Classical Quantum Gravity, 33 (2016), pp. 025003, 22.
- [2] G. S. ASANOV, *Finsler geometry, relativity and gauge theories*, Fundamental Theories of Physics, D. Reidel Publishing Co., Dordrecht, 1985.
- [3] J. K. BEEM, *Indefinite Finsler spaces and timelike spaces*, Canad. J. Math., 22 (1970), pp. 1035–1039.
- [4] M. A. JAVALOYES AND M. SÁNCHEZ, *Finsler metrics and relativistic spacetimes*, Int. J. Geom. Methods Mod. Phys., 11 (2014), p. 1460032 (15 pages).
- [5] M. A. JAVALOYES AND M. SÁNCHEZ, *Definition and examples of Finsler spacetimes*. Preprint, 2017.
- [6] V. A. KOSTELECKÝ, *Riemann-Finsler geometry and Lorentz-violating kinematics*, Phys. Lett. B, 701 (2011), pp. 137–143.
- [7] C. PFEIFER AND M. WOHLFARTH, *Causal structure and electrodynamics on finsler space-times*, Phys. Rev. D, 84 (2011).